



# Electro-Metabolic Coupling Investigated with Jitter Invariant Dictionary Learning

Sebastian Hitziger, Maureen Clerc, Alexandre Gramfort, Sandrine Sallet,  
Christian G. B  nar, Th  odore Papadopoulos

## ► To cite this version:

Sebastian Hitziger, Maureen Clerc, Alexandre Gramfort, Sandrine Sallet, Christian G. B  nar, et al..  
Electro-Metabolic Coupling Investigated with Jitter Invariant Dictionary Learning. OHBM 2014, Jun  
2014, Hamburg, Germany. . hal-01094674

**HAL Id: hal-01094674**

**<https://hal.inria.fr/hal-01094674>**

Submitted on 12 Dec 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin  e au d  p  t et    la diffusion de documents scientifiques de niveau recherche, publi  s ou non,   manant des   tablissements d'enseignement et de recherche fran  ais ou   trangers, des laboratoires publics ou priv  s.



# Electro-Metabolic Coupling Investigated with Jitter Invariant Dictionary Learning

Sebastian Hitziger<sup>1</sup>, Maureen Clerc<sup>1</sup>, Alexandre Gramfort<sup>2</sup>, Sandrine Saitlet<sup>3</sup>, Christian Bénar<sup>4</sup>, Théodore Papadopoulos<sup>1</sup>

<sup>1</sup> Project-Team Athena, Inria Sophia Antipolis, France

<sup>3</sup> Institut de Neurosciences des Systèmes, Marseille, France

sebastian.hitziger@inria.fr

<sup>2</sup> Telecom ParisTech - CEA Neurospin, Paris, France

<sup>4</sup> INSERM U1106, Marseille, France

<https://team.inria.fr/athena/>




This work aims at establishing a relationship between neurophysiological and hemodynamic activity in an animal model of epilepsy. For the analysis, we propose a novel algorithm that is

suited to learn meaningful representations of the multimodal datasets. As a result, we are able to learn a hemodynamic response and discover spike synchronization with hemodynamic activity.

## Goals and Challenges

### Goals:

- Investigate neurophysiological and hemodynamic activity.
- Establish relationship between the two.



Data: ~ 1 hour of simultaneous recordings of lead field potential (LFP) and laser doppler (LD) in Wistar-Kyoto rat. Bicuculline (a blocker of inhibition) was injected in the cortex to elicit epileptic-like discharges.

### Difficulties:

- Large dataset with several hundred epileptic spikes.
- Spikes change over time (shape, amplitude, spiking rate).
- Overlaps in spikes and hemodynamic responses.


## Processing LFP Data: Two Approaches

**A) Epoching:** LFP data is segmented into time windows centered around spikes [1] Drawbacks: (a) spikes need to be detected previously, (b) overlaps are not properly treated.

**B) Continuous model:** LFP data  $\mathbf{x}$  is composed of  $K$  spike shapes  $\{\mathbf{d}_i\}_{i=1}^K$  repeated at different latencies  $\delta_{ij}$  with varying amplitudes  $a_{ij}$  plus some noise  $\epsilon$ :

$$\mathbf{x} = \sum_{i=1}^K \sum_{j=1}^{M_i} a_{ij} \mathbf{d}_i(\cdot - \delta_{ij}) + \epsilon$$

Calculate  $a_{ij}$ ,  $\delta_{ij}$ ,  $\mathbf{d}_i$  using  $K$ -hierarchical alternate minimization:




Algorithm 1: Outline for learning waveforms (here spikes) and their latencies  $\delta_{ij}$  and amplitudes  $a_{ij}$ . Algorithm starts learning a single waveform  $\mathbf{d}_1$ , which is initialized with e.g. an arbitrary spike in the data (2). Then all spikes are detected and their latencies  $\delta_{ij}$  and amplitudes  $a_{ij}$  are calculated (3a, details below). The  $a_{ij}$ ,  $\delta_{ij}$  are now used to update waveforms  $\mathbf{d}_i$ . Steps 3a and 3b are repeated until convergence. If the residual still contains much structure, the previous steps may be repeated with an additional waveform.

### Core of Algorithm 1 - alternate minimization

- 3a** Correlation based detection of waveforms, can be solved by sparse coding (e.g. LARS [2]) with all possible shifts of the  $\mathbf{d}_i$ .
- 3b** Simple averaging if detected waveforms are not overlapping. Otherwise overlaps are accounted for in linear system (discrete deconvolution), which may be ill-conditioned for strong overlaps.

## Results on LFP


Algorithm 1 produces a compact representation of the data, providing additional insight.



Left: four spikes were learned on the data, note that their amplitudes are normalized. Their shapes differ mainly in duration. Right: amplitudes of detected spikes (top) are decreasing with time. They appear strongly negatively correlated with their local spiking rate (defined as  $1/\text{distance to previous spike [s]}$ ).

## LFP vs. Laser Doppler


**Hemodynamic response** is learned using Algorithm 1, where we take as  $\delta_{ij}$  the previously detected spike latencies.



Hemodynamic response learned on laser doppler using Algorithm 1 with  $K=1$  and  $\delta_{ij}$  the learned spike latencies. The presence of the oscillations with 7s period probably result from strong overlapping effects due to high spiking rates and long doppler response.

The figures show the laser doppler data (blue) and the approximation (red) given by the learned representation. During learning with Algorithm 1 we included a constant waveform  $\mathbf{d}_0$ , see green baseline. Note that relation between spike occurrence (red asterisk, bottom figures) and doppler response is only clearly visible during low spiking rates.

**Time-frequency analysis** reveals synchronization of spikes with 0.9-1 Hz rhythm in hemodynamics, probably related to respiration.



Local spiking rates ( $1/\text{distance [s]}$  to previous spike) are plotted on time-frequency map of laser doppler. During the three spike outbursts around 750, 1250, and 1900 seconds the spikes appear to synchronize with 0.9-1 Hz activity in the doppler. Around 750 seconds local spiking rates even clearly correspond to the subharmonics of this rhythm.

## Summary

### Proposed algorithm

- Produces informative data representations suited to analyze and compare multimodal data.
- Stepwise structure makes it flexible and allows integration of prior information.

### Electro-Metabolic Coupling

- Hemodynamic response was learned that well explains the data.
- Synchronization of spiking rates with periodic activity of 0.9-1 Hz in laser doppler becomes apparent in time-frequency map.